


Journal Pre-proof

Experience of different upper respiratory tract sampling strategies for detection of COVID-19

Guangming Ye, Yirong Li, Mengxin Lu, Song Chen, Yongwen Luo, Shichan Wang, Yejingpeng Wang, Xinghuan Wang



PII: S0195-6701(20)30111-0

DOI: <https://doi.org/10.1016/j.jhin.2020.03.012>

Reference: YJHIN 5940

To appear in: *Journal of Hospital Infection*

Received Date: 3 March 2020

Accepted Date: 6 March 2020

Please cite this article as: Ye G, Li Y, Lu M, Chen S, Luo Y, Wang S, Wang Y, Wang X, Experience of different upper respiratory tract sampling strategies for detection of COVID-19, *Journal of Hospital Infection*, <https://doi.org/10.1016/j.jhin.2020.03.012>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 The Healthcare Infection Society. Published by Elsevier Ltd. All rights reserved.

Experience of different upper respiratory tract sampling strategies for detection of COVID-19

Guangming Ye^{1,#}, Yirong Li^{1,#}, Mengxin Lu², Song Chen², Yongwen Luo², Shichan Wang¹, Yejingpeng Wang², Xinghuan Wang^{2,3,*}

¹Department of Clinical laboratory, Zhongnan Hospital of Wuhan University, Wuhan, 430072, China

²Department of Urology, Zhongnan Hospital of Wuhan University, Wuhan, 430072, China

³Center for Evidence-Based and Translational Medicine, Zhongnan Hospital of Wuhan University, Wuhan, 430072, China

[#]These authors contributed equally to this work.

***Corresponding author:** Dr. Xinghuan Wang, Email: wangxinghuan@whu.edu.cn, Tel. +86-27-6781-3104.

COVID-19 is spreading rapidly around the world¹. At present, the diagnosis of COVID-19 mainly depends on real-time RT-PCR assay of throat swabs². However, the false negative rate of nucleic acid test for SARS-CoV-2 with throat swabs is high, and throat swabs are uncomfortable for patients, and may induce coughing. The lingual swab is convenient and may achieve the same effect. This study compares the differences between lingual swab and throat swab sampling results, and analyzes whether standardized sampling by the same nurse could improve the detection rate compared with sampling by several nurses.

To analyze the positive rates of throat swabs and lingual swabs to detect COVID-19, we conducted a cohort study of two groups of suspected patients at two COVID-19 fixed-point hospitals (Wuhan No.7 Hospital and Zhongnan Hospital of Wuhan University). Zhongnan Hospital is one of the largest hospitals in the area, and has admitted a large number of COVID-19 patients^{3,4}. Two groups of consented patients were sampled using both sampling methods; all swabs were tested using a real-time reverse transcription-polymerase chain reaction (RT-PCR) assay for SARS-CoV-2. The first group contained 46 patients, who were from fever clinic of Wuhan No.7 Hospital (a COVID-19 fixed-point hospital). This group of patients was sampled by the same experienced, who used a standardized sampling method. Of these patients, 54.3% (25/46) were positive by RT-PCR; 54.3% (25/46) tested positive on throat swabs and 36.9% (17/46) were positive on lingual swabs; all patients with positive lingual swabs also had positive throat swabs. In another group, 45 patients from the fever clinic of Zhongnan Hospital of Wuhan University were also sampled by using both methods, but this group of patients was sampled by several nurses. 48.9% (22/45) were diagnosed with SARS-CoV-2 infection by RT-PCR. However this time 33.3% (15/45) tested positive on throat swabs and 35.6% (16/45) on lingual swabs; only 45.5%

(10/22) of the positive patients were detected by both methods (Table 1).

In total, of these 91 patients, the positive rate of throat swabs (44.0%) was higher than that of lingual swabs (36.3%), but this difference may have been attributable to a single experienced nurse collecting all of the samples in the first group (54.3% VS 36.9%). This suggests that highly trained or experienced nurses may improve the diagnostic sensitivity with throat swabs.

Furthermore, two Venn diagrams used to describe the relationship between two sampling methods in two groups (Supplementary Figure) illustrate the greater consistency of sampling in the first group, in which all lingual swab-positive patients also had a positive throat swab. The Kappa value was also higher (0.6599 VS 0.4592) for patients from Wuhan No.7 Hospital, sampled by the same experienced nurse, than for patients from Zhongnan Hospital (sampling performed by several nurses) (Supplementary Table).

In conclusion, the positive rate of throat swabs is higher than that of lingual swabs for the detection of COVID-19; however, in our small study this difference was only seen when swabs were collected by a single experienced nurse. When multiple nurses took samples, throat swabs had no obvious advantage over lingual swabs, and diagnostic sensitivity was improved by collection of samples from both sites. We believe that these observations should be of value to other centres establishing COVID-19 diagnostic programmes.

Conflict of interest statement

None declared.

Funding sources

This study was supported by the Medical Science Advancement Program (Clinical Medicine) of Wuhan University (TFLC2018002).

Journal Pre-proof

Reference

1. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* 2020; 382:727–33.
2. Chinese diagnosis and treatment plan of COVID-19 patients (The sixth edition). <http://www.nhc.gov.cn/yzygj/s7653p/202002/8334a8326dd94d329df351d7da8aefc2.shtml>.
3. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, *et al.* Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* 2020 Feb 7. doi: 10.1001/jama.2020.1585.
4. Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, *et al.* Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet* 2020 Jan 1. doi.org/10.1016/S0140-6736(20)30360-3.

Table 1. Positive rate of two sampling methods for nucleic acid detection in two hospitals.

	Throat swab positive cases/Total	Lingual swab positive cases/Total	Both positive cases/Total	Positive cases/Total	Both positive cases/ Positive cases
Wuhan No.7 Hospital					
Cases (n)	25/46	17/46	17/46	25/46	17/25
Rate (%)	54.3	36.9	36.9	54.3	68.0
Zhongnan Hospital					
Cases (n)	15/45	16/45	10/45	22/45	10/22
Rate (%)	33.3	35.6	22.2	48.9	45.5
Wuhan No.7 Hospital & Zhongnan Hospital					
Cases (n)	40/91	33/91	27/91	47/91	27/47
Rate (%)	44.0	36.3	29.7	51.6	57.4